

HK 1: Structure and Dynamics of Nuclei I

Time: Monday 15:00–16:30

Location: HS 2 Physik

Group Report

HK 1.1 Mon 15:00 HS 2 Physik
Systematic investigation of $E1$ strength below S_n in the tin isotopic chain using the $(d,p\gamma)$ reaction — ●MARKUS MÜLLENMEISTER, MICHAEL WEINERT, FLORIAN KLUWIG, TANJA SCHÜTTLER, and ANDREAS ZILGES — University of Cologne, Institute for Nuclear Physics, Germany

The evolution of the electric dipole ($E1$) response in various nuclei has been the subject of intense study for decades [1]. The structure of the so-called Pygmy Dipole Resonance, which emerges around and below the neutron separation energy of most medium- to heavy-mass nuclei, is of particular interest [2]. New insights into its origin can be gained through the well-established method of neutron transfer [3]. When combined with state-of-the-art analysis techniques, theoretical calculations, and comparative studies using different probes, the dipole strength can be examined across different nuclides to obtain detailed structural information [4].

In this contribution, a comparison for the tin isotopes, accessible via the $(d,p\gamma)$ -reaction, will be presented and compared to real photon scattering data. Together, these methods highlight the dominance of single-particle excitations at lower energies, while more complex configurations become significant at higher energies.

Supported by the DFG (ZI 510/10-1)

- [1] A. Bracco *et al.*, Prog. Part. Nucl. Phys. **106** (2019) 360
- [2] D. Savran *et al.*, Prog. Part. Nucl. Phys. **70** (2013) 210
- [3] M. Weinert *et al.*, Phys. Rev. Lett. **127** (2021) 242501
- [4] D. Savran *et al.*, Phys. Lett. B **786** (2018) 16

HK 1.2 Mon 15:30 HS 2 Physik
Photoabsorption Cross Sections of Tin and Calcium Isotopes — ●MARTIN BAUMANN¹, THOMAS AUMANN^{1,2}, MAIKE BEUSCHLEIN¹, ISABELLE BRANDHERM¹, MEY TAL DUER¹, AMRITA GUPTA¹, PHILLIP IMGRAM¹, ANDREA JEDELE¹, LIANCHENG JI¹, IGOR JUROSEVIC¹, MARCO KNÖSEL¹, NIKOLINA WAGNER¹, ENIS LORENZ¹, HANNES MAYR¹, LEANDRO MILHOMES DA FONSECA¹, NIKHIL MOZUMDAR¹, ANN ROCHELE NETTO¹, OLIVER PAPST¹, THOMAS POHL¹, HEIKO SCHEIT¹, GERHART STEINHILBER¹, SONJA STORCK-DUTINE¹, DMYTRO SYMOCHKO¹, IYABO USMAN³, and PATRICK VAN BEEK¹ — ¹Institut für Kernphysik, TU Darmstadt, Germany — ²GS

Helmholtzzentrum, Darmstadt, Germany — ³University of the Witwatersrand, Johannesburg, South Africa

The photoabsorption setup of the NEPTUN photon tagger at the superconducting linear accelerator S-DALINAC has been used to investigate the photoabsorption cross sections of Sn-112,116,120,124 as well as Ca-40,48 in the region from 5 to 30 MeV using a beam of tagged photons. In this talk the measurement method as well as the current status of the data analysis will be presented.

This work is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID 279384907 - SFB 1245.

HK 1.3 Mon 15:45 HS 2 Physik
Probing the doubly-magic shell closure at ^{132}Sn by Coulomb excitation of neutron-rich ^{130}Sn — ●MAXIMILIAN DROSTE¹, PETER REITER², and THORSTEN KRÖLL² for the IS702-Collaboration — ¹IKP, Universität zu Köln — ²IKP, TU Darmstadt

Excited states of ^{130}Sn , the even-even neighbour of doubly-magic ^{132}Sn , were studied by safe Coulomb excitation using the highly-efficient MINIBALL array. The ^{130}Sn ions were accelerated to 4.4 MeV/u at the HIE-ISOLDE accelerator and collided with a ^{206}Pb target. Deexciting γ rays from excited states were detected in coincidence with scattered particles. In addition to γ rays from the first 2^+ state of ^{130}Sn , deexcitation from higher-lying states was observed, attributed to an isomeric $^{130}\text{Sn } J^\pi = 7^-$ beam component. A new $B(E2; 0_{g.s.}^+ \rightarrow 2_1^+)$ value is compared to recent theoretical results from

state-of-the-art MCSM calculations which differ strongly from previous measurements [1,2]. These calculations also indicate a transition from a slightly oblate to a prolate configuration of the first excited 2^+ state across doubly magic ^{132}Sn . The high statistics of the performed experiment allows for an experimental investigation of the Q_{2^+} value.

[1] D. Rosiak *et al.* Phys. Rev. Lett. **121**, 252501 (2018)

[2] T. Togashi *et al.* Phys. Rev. Lett. **121**, 062501 (2018)

Supported by the German BMBF 05P21PKCI1, 05P24PKCI1, 05P21RDCI2 and European Union*s Horizon Europe Framework research and innovation programme under grant agreement no. 101057511

HK 1.4 Mon 16:00 HS 2 Physik
Lifetimes of excited states in $^{116,118}\text{Sn}$ — ●SARAH PRILL, ELIAS BINGER, ANNA BOHN, TOBIAS LANGEL, MICHAEL WEINERT, and ANDREAS ZILGES — University of Cologne, Institute for Nuclear Physics, Germany

The proton-magic tin isotopic chain has the highest number of stable isotopes and is therefore an ideal candidate to study nuclear properties in a wide range of nuclei. Lifetimes of excited states in the femtosecond range have already been determined in Cologne for ^{112}Sn and ^{114}Sn [1] using the Coincidence Doppler-Shift Attenuation Method (CDSAM) [2,3]. The study was continued for the nuclei ^{116}Sn and ^{118}Sn , using not only inelastic proton scattering but also alpha particles as the beam.

Coincidence data were recorded with the SONIC@HORUS setup [4] at the tandem accelerator of the University of Cologne. Combining particle and γ -ray detection allowed the reconstruction of the reaction kinematics and enabled the analysis of single levels without feeding contributions.

This presentation will show the lifetime results for ^{116}Sn and ^{118}Sn and discuss the influence of different beam particles.

Supported by the DFG (ZI 510/9-2).

[1] M. Spieker *et al.*, Phys. Rev. C **97**, 054319 (2018).

[2] A. Hennig *et al.*, Nucl. Instr. Meth. A **758**, 171 (2015).

[3] S. Prill *et al.*, Phys. Rev. C **105**, 034319 (2022).

[4] S. G. Pickstone *et al.*, Nucl. Instr. Meth. A **875**, 104 (2017).

HK 1.5 Mon 16:15 HS 2 Physik

Determination of the energy-resolvable $E1$ - and $M1$ -strength distribution in ^{70}Zn — ●J. HAUF¹, V. WERNER¹, A. D. AYANGEAKAA^{2,3}, D. BALABANSKI^{4,5}, M. BEUSCHLEIN¹, R. BEYER⁶, S. W. FINCH^{2,7}, A. GUPTA¹, D. GRIBBLE^{2,3}, T. HENSEL⁶, M. HEUMÜLLER¹, F. E. IDOKU^{2,3}, J. ISAAK¹, X. JAMES^{2,3}, R. V. F. JANSSENS^{2,3}, S. R. JOHNSON^{2,3}, A. JUNGHANS⁶, J. KLEEMANN¹, P. KOSEOGLU¹, T. KOWALEWSKI^{2,3}, A. KUSOGLU⁴, J. LU¹, E. MASHA⁶, C. M. NICKEL¹, O. PAPST¹, M. PICHOTTA⁶, N. PIETRALLA¹, K. PRIFTI¹, K. RÖMER⁶, A. SARACINO^{2,3}, P.-A. SÖDERSTRÖM⁴, K. SCHMIDT⁶, R. SCHWENGER⁶, A. THEES⁶, S. TURKAT⁶, J. VOGEL¹, A. WAGNER⁶, and A. YADEV⁶ — ¹TU Darmstadt, IKP — ²TUNL — ³University of North Carolina — ⁴ELI-NP — ⁵Horia Hulubei National Institute — ⁶Helmholtz-Zentrum Dresden-Rossendorf — ⁷Duke University

Nuclear resonance fluorescence experiments with bremsstrahlung and quasi-monoenergetic photon beams have been conducted on ^{70}Zn at γELBE and $\text{HI}\gamma\text{S}$. The investigation of the most neutron-rich stable zinc isotope aims to achieve a better understanding of nuclear structure phenomena, such as shape coexistence and the Pygmy dipole resonance, at the $N = 40$ harmonic oscillator shell closure. The status of the analysis, including spectra and preliminary results for the $E1$ - and $M1$ -strength distributions, are shown and discussed.

This work is supported by DFG under Project-IDs 499256822*GRK 2891, 279384907*SFB 1245 and by ELI-RO under ELI-RO/RDI/2024 002 and ELI-RO/RDI/2024 007.